



# Medical Coverage Policy

Effective Date .....5/15/2025

Next Review Date .....5/15/2026

Coverage Policy Number..... 0073

## Cardiac Rehabilitation (Phase II Outpatient)

### Table of Contents

Overview ..... 2  
 Coverage Policy ..... 2  
 Health Equity Considerations..... 2  
 General Background..... 3  
 Medicare Coverage Determinations ..... 13  
 Coding Information ..... 13  
 References..... 14  
 Revision Details ..... 22

### Related Coverage Resources

- [Heart, Lung, and Heart-Lung Transplantation](#)
- [Transcatheter Heart Valve Procedures](#)
- [Ventricular Assist Devices \(VADs\),](#)
- [Percutaneous Cardiac Support Systems and](#)
- [Total Artificial Heart](#)

### INSTRUCTIONS FOR USE

*The following Coverage Policy applies to health benefit plans administered by Cigna Companies. Certain Cigna Companies and/or lines of business only provide utilization review services to clients and do not make coverage determinations. References to standard benefit plan language and coverage determinations do not apply to those clients. Coverage Policies are intended to provide guidance in interpreting certain standard benefit plans administered by Cigna Companies. Please note, the terms of a customer’s particular benefit plan document [Group Service Agreement, Evidence of Coverage, Certificate of Coverage, Summary Plan Description (SPD) or similar plan document] may differ significantly from the standard benefit plans upon which these Coverage Policies are based. For example, a customer’s benefit plan document may contain a specific exclusion related to a topic addressed in a Coverage Policy. In the event of a conflict, a customer’s benefit plan document always supersedes the information in the Coverage Policies. In the absence of a controlling federal or state coverage mandate, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of 1) the terms of the applicable benefit plan document in effect on the date of service; 2) any applicable laws/regulations; 3) any relevant collateral source materials including Coverage Policies and; 4) the specific facts of the particular situation. Each coverage request should be reviewed on its own merits. Medical directors are expected to exercise clinical judgment where appropriate and have discretion in making individual coverage determinations. Where coverage for care or services does not depend on specific circumstances, reimbursement will only be provided if a requested service(s) is submitted in accordance with the relevant criteria outlined in the applicable Coverage Policy, including covered diagnosis and/or procedure code(s). Reimbursement is not allowed for services when billed for conditions or diagnoses that are not covered under this Coverage Policy (see “Coding Information” below). When billing, providers must use the most appropriate codes as of the effective date of the submission. Claims submitted for services that are not accompanied by covered code(s) under the applicable Coverage Policy*

*will be denied as not covered. Coverage Policies relate exclusively to the administration of health benefit plans. Coverage Policies are not recommendations for treatment and should never be used as treatment guidelines. In certain markets, delegated vendor guidelines may be used to support medical necessity and other coverage determinations.*

## Overview

This Coverage Policy addresses cardiac rehabilitation (Phase II) services that are provided on an outpatient basis post facility discharge.

## Coverage Policy

**Coverage for cardiac rehabilitation (CR) varies across plans. Refer to the customer's benefit plan document for coverage details.**

**If benefit coverage is available for cardiac rehabilitation, then the following conditions apply.**

**A medically supervised outpatient Phase II Cardiac Rehabilitation program (CPT®\* code 93797, 93798) is considered medically necessary following ANY of the following events:**

- acute myocardial infarction (MI)
- coronary artery bypass grafting (CABG)
- percutaneous coronary vessel remodeling
- valve replacement or repair
- coronary artery disease (CAD) associated with chronic stable angina that has failed to respond adequately to pharmacotherapy and is interfering with the ability to perform age-related activities of daily living and/or impairing functional abilities
- heart failure that has failed to respond adequately to pharmacotherapy and is interfering with the ability to perform age-related activities of daily living and/or impairing functional abilities
- following surgical septal myectomy via thoracotomy
- heart transplantation or heart-lung transplantation
- major pulmonary surgery, great vessel surgery, or MAZE arrhythmia surgery
- placement of a ventricular assist device
- sustained ventricular tachycardia or fibrillation
- survivors of sudden cardiac arrest

**EACH of the following is considered educational and/or training in nature and not medically necessary:**

- phase III or IV cardiac rehabilitation programs
- intensive cardiac rehabilitation programs (HCPCS code G0422, G0423) (e.g. Pritikin Program, Ornish Program for Reversing Heart Disease, Benson-Henry Institute Cardiac Wellness Program)

## Health Equity Considerations

Health equity is the highest level of health for all people; health inequity is the avoidable difference in health status or distribution of health resources due to the social conditions in which people are born, grow, live, work, and age.

Social determinants of health are the conditions in the environment that affect a wide range of health, functioning, and quality of life outcomes and risks. Examples include safe housing, transportation and neighborhoods; racism, discrimination and violence; education, job opportunities and income; access to nutritious foods and physical activity opportunities; access to clean air and water; and language and literacy skills.

Socioeconomic status as measured by several individual (e.g., income, education attainment, occupation, medical assistance such as Medicaid) and environmental indicators (e.g., area deprivation index or median income based on zip code data) has been shown to have a significant impact on cardiovascular disease (CVD) development and outcomes. Individuals with low socioeconomic status (SES) and CVD face a disproportionately higher risk of recurrent events and mortality compared to those with high SES.

In spite of the known benefits, cardiac rehabilitation (CR) programs are consistently underutilized. It is estimated that less than 40% of eligible patients enroll in CR after a qualifying event. A major factor is the under-referral of patients to CR, especially women, older adults, and under-represented racial and ethnic groups. According to an analysis by Oehler et al. (2023), those living in a rural setting are also more likely to underutilize cardiac rehabilitation services. Referral rates for women are approximately 10% lower than for men, even after adjusting for age and comorbid conditions. Lack of transportation, lack of enjoyment, and home responsibilities are cited more frequently by women as reasons for nonparticipation.

Black patients are referred 20% less frequently than are non-Hispanic white patients. Enrollment is also low, as Black patients participate in CR half as often as white patients (17.3% vs. 30%). These findings are particularly concerning, as women and non-whites are significantly more likely to die within five years after a first myocardial infarction, compared with white male patients. One proposed solution is for hospitals to implement an automated referral process, to prevent referral bias (Mathews and Brewer, 2021; Balady, et al., 2011).

Rural patients face enrollment challenges because of limited access. An American Heart Association (AHA) presidential advisory named poor rural access to phase II CR as a contributing factor to poor cardiovascular outcomes. Those living more than 15 miles from a CR center are 71% less likely to be enrolled than are those who live less than 1.5 miles away. Additionally, virtual models might be problematic in rural areas because of limited broadband internet access (24% of households do not have broadband internet in urban settings vs. 32% of households in rural settings).

Finally, there is a gradual drop in participation in CR as patients age, dropping to one-third of baseline values in those over 85 years old. Older age and functional and sensory impairments common in aging may lead to lower rates of participation. Virtual, home-based interventions suggest benefit.

## General Background

### Cardiac Rehabilitation

The 2005 American Heart Association/American Association of Cardiovascular and Pulmonary Rehabilitation (AHA/AACVPR) scientific statement defines cardiac rehabilitation (CR) as coordinated, multifaceted interventions designed to optimize a cardiac patient's physical,

psychological, and social functioning, in addition to stabilizing, slowing, or even reversing the progression of the underlying atherosclerotic processes, thereby reducing morbidity and mortality (Leon, et al., 2005). CR typically incorporates exercise training, patient education, and health behavior modification to improve outcomes in individuals with cardiovascular disease.

The candidates for CR/secondary prevention programs are patients who recently have had a myocardial infarction (MI); have undergone coronary artery bypass graft surgery (CABG) or percutaneous coronary interventions; heart transplant candidates or recipients; or patients with stable chronic heart failure, peripheral arterial disease with claudication, or other forms of cardiovascular disease or cardiac surgical procedures (Leon, et al., 2005).

CR/secondary prevention programs currently include baseline patient assessments, nutritional counseling, aggressive risk-factor management (i.e., lipids, hypertension, weight, diabetes, and smoking), psychosocial and vocational counseling, and physical activity counseling and exercise training. Additionally, CR programs include the appropriate use of cardioprotective drugs that have evidence-based efficacy for secondary prevention (Leon, et al., 2005).

The early CR programs initiated mobilization after a myocardial infarction and were referred to as Phase I or inpatient CR. The goal was to condition the patient to safely carry out activities of daily living following discharge. Such programs entailed prescribing activity in rigid steps with successively higher metabolic equivalents (METs). Comprehensive CR programs eventually grew to include three to four phases.

- **Phase I (Inpatient):** Inpatient rehabilitation, usually lasting for the duration of hospitalization for an acute coronary event or surgery. It emphasizes a gradual, progressive approach to exercise and an education program that helps the patient understand the disease process, the rehabilitation process, and initial preventive efforts to slow the progression of disease. Submaximal exercise testing before hospital discharge is done to provide important prognostic information and help restore patient confidence. These programs are uncommon due to the brevity of most hospital stays.
- **Phase II (Outpatient Medically Supervised):** Multifaceted, physician-directed outpatient rehabilitation, lasting from hospital discharge to 2–12 weeks later. Phase II CR emphasizes safe physical activity to improve conditioning with continued behavior modification aimed at smoking cessation, weight loss, healthy eating, and other factors to reduce disease risk.
- **Phase III (Supervised, Transitional):** Supervised rehabilitation, often in a group setting, lasting 6–12 months. Establishes a prescription for safe exercise that can be performed at home or in a community service facility, such as a senior center, and continues to emphasize risk-factor reduction while transitioning to independence.
- **Phase IV (Maintenance/Follow-Up):** This is usually an indefinite program, and some programs may combine Phases III and IV. The goal is to encourage lifelong adherence to the healthy habits established during Phase II. Follow-up visits can occur at 6–12 month intervals. Blood pressure and pulse measurement, serum lipid levels, and even repeat maximal exercise tolerance tests can provide useful feedback to the patient and indicate areas that may require lifestyle changes to minimize coronary events.

**Phase II (Outpatient) Cardiac Rehabilitation (CR):** Phase II CR is described by the U.S. Public Health Service as consisting of “comprehensive, long term programs involving medical evaluation, prescribed exercise, cardiac risk factor modification, education, and counseling”. These programs “are designed to limit the physiologic and psychological effects of cardiac illness, reduce

the risk of sudden death or reinfarction, control cardiac symptoms, stabilize or reverse the atherosclerotic process, and enhance the psychosocial and vocational status of selected patients.” CR programs aim to reduce subsequent cardiovascular related morbidity and mortality. Phase II CR specifically refers to outpatient, medically supervised programs that provide both electrocardiogram (ECG) monitored and non-electrocardiogram (ECG) monitored sessions. The programs are typically initiated within one to three weeks after hospital discharge and generally administered within the six months following discharge from the hospital (Bartels, et al., 2024).

It is recommended that patients referred to CR undergo a symptom-limited exercise tolerance/stress test before entering the CR program. The exercise test is to exclude important symptoms, ischemia, or arrhythmias that might require other interventions before exercise training. The exercise test also serves to establish baseline exercise capacity and to determine maximum heart rate for use in preparing an exercise prescription. These tests are generally done with the patient on their usual medications to mimic the heart rate response likely to occur during exercise training. Exercise intensity is regulated by monitoring peak heart rate. The exercise training modalities used during Phase II, as in Phase I, usually consist of walking and stationary bicycling, and the patient and family are educated about coronary risk and self-monitoring.

Most Phase II exercise programs consist of three sessions per week for 12 weeks, however the frequency and duration may be impacted by the level of cardiac risk stratification. The CR program is individualized by assessing the patient’s history and current need for cardiac risk factor modification. Risk stratification is used to identify patients at risk for death or reinfarction, and to provide guidelines for the rehabilitative process.

Each cardiac rehabilitation session is individualized to meet the patient’s needs. Exercise training is the principal component of the program, as it results in increased peak exercise capacity, usually expressed in METs. The MET is the total oxygen requirement of the body, with one MET equal to 3.5 milliliters of oxygen consumed per kilogram of body weight per minute. Exercise training is aimed to improve MET capacity, resulting in improved oxygen delivery and extraction, by exercising skeletal muscles, decreasing the cardiovascular requirements of exercise and increasing the amount of work that can be done before ischemia (i.e., blood deficiency) occurs.

Contraindications to the exercise program component of CR include the following (Davis, 2019):

- unstable angina
- resting systolic blood pressure >200 mm Hg or diastolic BP >100 mm Hg
- orthostatic blood pressure drop or drop during exercise training of >20 mm Hg
- third-degree heart block
- resting ST displacement (> 3 mm)
- uncontrolled diabetes
- acute systemic illness or fever
- recent embolism
- active pericarditis or myocarditis
- moderate to severe aortic stenosis
- thrombophlebitis
- uncontrolled arrhythmias
- uncontrolled congestive heart failure (CHF)
- orthopedic problems that prohibit exercise

**Centers for Medicare and Medicaid Services (CMS):** CMS currently covers CR for the following indications (CMS, 2024):

- a documented acute myocardial infarction (AMI) within the preceding 12 months

- CABG surgery
- stable angina pectoris
- heart valve replacement/repair
- percutaneous transluminal coronary angioplasty (PTCA) or coronary artery stenting
- heart or heart/lung transplant
- stable, chronic heart failure (defined as patients with left ventricular ejection fraction of 35% or less and New York Heart Association (NYHA) class II to IV symptoms despite being on optimal heart failure therapy for at least 6 weeks)
- other cardiac conditions as specified through a national coverage determination (NCD)

CMS lists the following cardiac rehabilitation program requirements:

- Physician-prescribed exercise each day cardiac rehabilitation items and services are furnished.
- Cardiac risk factor modification, including education, counseling, and behavioral intervention at least once during the program, tailored to individual needs.
- Psychosocial assessment; outcomes assessment; and an individualized treatment plan detailing how components are utilized for each individual.
- Outcomes assessment.
- An individualized treatment plan detailing how components are utilized for each patient, established, reviewed, and signed by a physician every 30 days.

In 2010, CMS updated criteria on the frequency and duration of cardiac rehabilitation services stating that cardiac rehabilitation items and services must be furnished in a physician's office or a hospital outpatient setting. All settings must have a physician immediately available and accessible for medical consultations and emergencies at all times items and services are being furnished under the program. Cardiac rehabilitation program sessions are limited to a maximum of two 1-hour sessions per day for up to 36 sessions over up to 36 weeks, with the option for an additional 36 sessions over an extended period of time if approved by the Medicare contractor.

**Literature Review:** Several Cochrane systematic reviews and meta-analyses of randomized controlled trials have evaluated the effectiveness of center-based cardiac rehabilitation (CR) for numerous indications, including heart failure, post-heart transplantation, post-myocardial infarction, and after percutaneous coronary intervention (PCI). Overall, the evidentiary analyses have concluded that CR, particularly exercise-based CR, confers clinically important improvements in exercise capacity and quality of life, and lowers the risk of rehospitalization and death (Buckley, et al., 2024; Molloy, et al., 2024; Gore, et al., 2023; Abraham, et al., 2021; Dibben, et al., 2021; Nielsen, et al., 2019; Long, et al., 2018; Anderson, et al., 2017a; Anderson and Taylor, 2014).

Clark et al. (2005), from the University of Alberta Evidence-based Practice Center for the AHRQ Technology Assessment Program, conducted a meta-analysis of coronary heart disease management programs. The purpose of the study was to determine the effectiveness of secondary cardiac prevention programs with and without exercise components. The interventions tested in the trials, and frequency and duration of the interventions, varied substantially among the studies. The studies enrolled highly selected patient populations. After reviewing 63 randomized controlled trials of 21,295 patients with coronary disease, the authors concluded that secondary prevention programs for patients already diagnosed with cardiac disease improved processes of care, enhanced quality of life/function status, reduced recurrent myocardial infarctions, reduced hospitalizations, and reduced long-term mortality in patients with established CAD.

**Professional Societies/Organizations:** Evidence-based professional society guidelines consistently and strongly recommend comprehensive center-based cardiac rehabilitation (CR) in the management and prevention of cardiovascular disease.

In developing the guidelines below, the American College of Cardiology (ACC)/American Heart Association (AHA) guideline task force used evidence-based methodologies to assign each recommendation a Class of Recommendation and a Level of Evidence. The Class of Recommendation indicates the degree of benefit versus risk and corresponds to the strength of the recommendation. The Level of Evidence indicates the certainty of the evidence supporting the recommendation; based on the type, size, quality, and consistency of the evidence reviewed. The class and evidence levels were updated in 2015 and 2019 to further refine the definitions and better reflect the evidence upon which the recommendation is based (Lawton, et al., 2022; Halperin, et al., 2016; O’Gara, et al., 2013).

<b>Class (Strength) of Recommendation</b>	
Guidelines published prior to August 2015	Guidelines published after August 2015
<p><b>Class I</b></p> <ul style="list-style-type: none"> <li>• Benefit &gt;&gt;&gt; Risk</li> <li>• Procedure/Treatment should be performed/administered.</li> </ul>	<p><b>Class 1 (Strong)</b></p> <ul style="list-style-type: none"> <li>• Benefit &gt;&gt;&gt; Risk</li> <li>• Intervention is recommended; is indicated/useful/effective/beneficial</li> </ul>
<p><b>Class IIa</b></p> <ul style="list-style-type: none"> <li>• Benefit &gt;&gt; Risk</li> <li>• Additional studies with focused objectives needed</li> <li>• It is reasonable to perform procedure/ administer treatment</li> </ul>	<p><b>Class 2a (Moderate)</b></p> <ul style="list-style-type: none"> <li>• Benefit &gt;&gt; Risk</li> <li>• Intervention is reasonable; can be useful/effective/beneficial</li> </ul>
<p><b>Class IIb</b></p> <ul style="list-style-type: none"> <li>• Benefit ≥ Risk</li> <li>• Additional studies with broad objectives needed; additional registry data would be helpful</li> <li>• Procedure/treatment may be considered</li> </ul>	<p><b>Class 2b (Weak)</b></p> <ul style="list-style-type: none"> <li>• Benefit ≥ Risk</li> <li>• Intervention may be reasonable; may be considered; its usefulness/ effectiveness is unknown/unclear/ uncertain or not well-established</li> </ul>
<p><b>Class III</b></p> <ul style="list-style-type: none"> <li>• Risk ≥ Benefit</li> <li>• Procedure/treatment should not be performed/administered, since it is not helpful and may be harmful</li> </ul>	<p><b>Class 3: No Benefit (Moderate)</b></p> <ul style="list-style-type: none"> <li>• Benefit = Risk</li> <li>• Intervention is not recommended/ indicated/useful/effective/beneficial; it should not be performed/administered</li> </ul>
	<p><b>Class 3: Harm (Strong)</b></p> <ul style="list-style-type: none"> <li>• Risk &gt; Benefit</li> <li>• Intervention is potentially harmful; causes harm; is associated with excess morbidity/mortality; should not be performed/administered</li> </ul>

<b>Level of Evidence (LOE)</b>	
Guidelines published prior to August 2015	Guidelines published after August 2015
<b>Level A</b> <ul style="list-style-type: none"> <li>Multiple populations evaluated</li> <li>Data derived from multiple randomized clinical trials (RCTs) or meta-analyses</li> </ul>	<b>Level A</b> <ul style="list-style-type: none"> <li>High-quality evidence from more than 1 RCT</li> <li>Meta-analyses of high-quality RCTs</li> <li>One or more RCTs corroborated by high-quality registry studies</li> </ul>
<b>Level B</b> <ul style="list-style-type: none"> <li>Limited populations evaluated</li> <li>Data derived from a single randomized trial or nonrandomized studies</li> </ul>	<b>Level B-R (Randomized)</b> <ul style="list-style-type: none"> <li>Moderate-quality evidence from 1 or more RCTs</li> <li>Meta-analyses of moderate-quality RCTs</li> </ul>
	<b>Level B-NR (Nonrandomized)</b> <ul style="list-style-type: none"> <li>Moderate-quality evidence from 1 or more well-designed, well-executed nonrandomized studies, observational studies, or registry studies</li> <li>Meta-analyses of such studies</li> </ul>
<b>Level C</b> <ul style="list-style-type: none"> <li>Very limited populations evaluated</li> <li>Only consensus opinion of experts, case studies, or standard of care</li> </ul>	<b>Level C-LD (Limited Data)</b> <ul style="list-style-type: none"> <li>Randomized or nonrandomized observational or registry studies with design or execution limitations</li> <li>Meta-analyses of such studies</li> <li>Physiological or mechanistic studies in human subjects</li> </ul>
	<b>Level C-EO (Expert Opinion)</b> <ul style="list-style-type: none"> <li>Consensus of expert opinion based on clinical experience</li> </ul>

The 2025 ACC/AHA/American College of Emergency Physicians (ACEP)/National Association of EMS Physicians (NAEMSP)/ Society for Cardiovascular Angiography and Interventions (SCAI) guideline for the management of patients with acute coronary syndromes (ACS) provided the following recommendations (Rao, et al., 2025):

- patients with ACS should be referred to an outpatient CR program prior to hospital discharge to reduce death, myocardial infarction, hospital readmissions, and improve functional status and quality of life (Class 1; LOE A)
- in patients with ACS, a home-based CR program is a reasonable alternative to a center-based CR program to improve functional status and quality of life (Class 2a; LOE B-R)

In 2024, the AHA and the American Association of Cardiovascular and Pulmonary Rehabilitation (AACVPR) updated their scientific statement addressing the core components of CR/secondary prevention programs. The update presented the current information on the evaluation, interventions, and expected outcomes in each of the core components of CR/secondary prevention programs, including:

- baseline patient assessment

- nutritional counseling
- cardiovascular disease and risk factor management:
  - dyslipidemia management
  - hypertension management
  - diabetes management
  - tobacco cessation
- weight management and body composition
- psychosocial management
- physical activity counseling
- aerobic exercise training
- strength training

The updated statement addressed the use of varied delivery methods for CR, including traditional center-based programs, virtual, remote, and hybrid CR program models. These newer care models may overcome barriers to accessing traditional CR programs (see next section), however the authors caution that “the adoption of novel models of delivery must not change the fundamental therapeutic foundation in what is delivered. CR is a medically supervised secondary prevention program with a physician medical director, day-to-day medical supervision by a physician or nonphysician practitioner, and core components, all of which must be included for a program to be classified as CR. Alternative delivery models that do not meet these statutorily defined standards are not CR and should not be viewed as an acceptable alternative to CR” (Brown, et al., 2024).

The AHA/AACVPR statement reconfirmed that aerobic exercise supplemented with strength training remains the cornerstone intervention for CR. Graded exercise testing was recommended as part of the initial assessment to help individualize the exercise intensity prescription. When graded exercise testing is not available, the Talk Test or 6-minute walk test may be used. Regarding continuous ECG monitoring during exercise interventions, the authors note that “continuous electrocardiographic telemetry and exercise blood pressure monitoring likely add value to a CR program, especially in high-risk or symptomatic patients. However, exercise is safe in general, and evidence is insufficient to determine how to best implement these services, how long to continue them, or which patients are most likely to benefit” (Brown, et al., 2024).

The 2022 AHA/ACC/Heart Failure Society of America (HFSA) guideline for the management of heart failure recommends exercise training (or regular physical activity) for patients with heart failure who are able to participate, to improve functional status, exercise performance, and quality of life (Class 1; Level of Evidence: A). Further, in patients with heart failure, a cardiac rehabilitation program can be useful to improve functional capacity, exercise tolerance, and health-related quality of life (Class 2a; Level of Evidence: B-NR) (Heidenreich, et al., 2022).

The 2021 ACC/AHA/SCAI guideline for coronary artery revascularization recommends that patients who have undergone revascularization should be prescribed a comprehensive cardiac rehabilitation program prior to hospital discharge or at the first outpatient visit, to reduce deaths and hospital readmissions and improve quality of life (Class: 1; Level of Evidence: A). Further, the guideline recommends that these patients should be educated about cardiovascular disease risk factors and how to modify them to reduce cardiovascular events (Class: 1; Level of Evidence: C-LD) (Lawton, et al., 2022).

The ACC/AHA clinical performance and quality measures for CR were updated most recently in 2018. The 2018 document retires the original “Set B” measures while publishing six new performance measures and three quality measures. These measures focus on the opportunities to improve referrals to outpatient CR from both inpatient and outpatient presentations. The updated performance measures state all patients hospitalized and evaluated in outpatient setting with a primary diagnosis of an acute myocardial infarction (MI) or chronic stable angina (CSA), or who during hospitalization have undergone coronary artery bypass graft (CABG) surgery, a

percutaneous coronary intervention (PCI), cardiac valve surgery, or cardiac transplantation are to be referred to an early outpatient cardiac rehabilitation/secondary prevention (CR) program. Furthermore, the performance measures state all patients evaluated in the inpatient and outpatient setting who within the past 12 months have a primary diagnosis of heart failure with reduced ejection fraction be referred to outpatient exercise training, typically delivered in an outpatient CR program. The remaining performance measures and quality measures focus on enrollment, adherence and clinical outcomes of the CR program. The authors noted that improved clinical outcomes are realized with a "full dose" of 36 prescribed sessions. CR communication to healthcare providers is important and care coordination is considered standard of care. The patients who are appropriate for entry into a CR program include persons 18 years of age or older who, during the previous year, have had one or more of the qualifying diagnoses previously noted. (Thomas, et al., 2007, 2010, 2018).

The updated 2011 AHA/ACCF secondary prevention and risk reduction therapy for patients with coronary and other atherosclerotic vascular disease guideline recommendations for cardiac rehabilitation states:

- All eligible patients with acute coronary syndrome (ACS) or whose status is immediately post coronary artery bypass surgery or post-PCI should be referred to a comprehensive outpatient cardiovascular rehabilitation program either prior to hospital discharge or during the first follow-up office visit (Class I; Level of Evidence: A).
- All eligible outpatients with the diagnosis of ACS, coronary artery bypass surgery or PCI (Class I; Level of Evidence: A), chronic angina (Class I; Level of Evidence: B), and/or peripheral artery disease (Class I; Level of Evidence: A) within the past year should be referred to a comprehensive outpatient cardiovascular rehabilitation program.
- A home-based cardiac rehabilitation program can be substituted for a supervised, center-based program for low-risk patients (Class I; Level of Evidence: A).
- A comprehensive exercise-based outpatient cardiac rehabilitation program can be safe and beneficial for clinically stable outpatients with a history of heart failure (Class IIa; Level of Evidence: B) (Smith, et al., 2011).

The 2011 AHA guidelines for the prevention of cardiovascular disease (CVD) in women indicated that a comprehensive CVD risk-reduction program (e.g., cardiovascular or stroke rehabilitation; physician-guided home- or community-based exercise training program) should be recommended to women with a recent acute coronary syndrome or coronary revascularization, new-onset or chronic angina, recent cerebrovascular event, peripheral arterial disease (Class I; Level of Evidence: A) or current/prior symptoms of heart failure and an LVEF  $\leq$  35% (Class I; Level of Evidence B) (Mosca, et al., 2011).

### **Virtual/Remote, Home-Based and/or Hybrid Cardiac Rehabilitation**

In recent years, an increasing number of alternative strategy programs have emerged to combat the low utilization of center-based CR, including home- or community-based ("remote"), virtual, and hybrid programs. These alternative programs vary in structure, length, and implementation, but generally rely on remote coaching with indirect exercise supervision that occurs outside of the traditional outpatient center or office setting. Technologies range from virtual, app-based programs that allow for real-time, two-way audiovisual communication between the individual and rehabilitation staff, to more remote programs which utilize only intermittent interaction via phone, email, and/or mail.

The proposed advantages of such programs include convenience and flexibility, which purportedly increase enrollment and adherence, especially among individuals who otherwise would be unable or unwilling to attend a center-based program. Barriers to implementing such programs include health data and privacy concerns; inequitable access to the required technology and/or broadband

service; variable technology literacy; depersonalization of the patient-provider relationship; reliance on self-reported outcomes; and unknown accuracy and reliability of digital technologies (e.g., wearable biosensors). Nonetheless, published evidence generally supports the safety and efficacy of virtual, home-based and hybrid CR programs. Further, several professional societies support the use of these alternate CR modalities, particularly by those individuals who are unable or unwilling to attend a center-based CR program (Brown, et al., 2024; Maddox, et al., 2024; Golbus, et al., 2023; Ghisi, et al., 2022; Thomas, et al., 2019).

**Literature Review:** Evidence in the peer-reviewed published scientific literature comparing home-based CR to no formal CR (i.e., medical management only), and/or to traditional, center-based CR programs consists of randomized controlled trials, prospective noncomparative trials, retrospective reviews, and meta-analyses. Most trials involve short-term follow-up (i.e.,  $\leq 12$  months), and vary widely in terms of CR program elements, program length, control groups, inclusion/exclusion criteria, outcome measures, and data reporting. Generally, studies have found that, compared to individuals who did not participate in a formal CR program, individuals in home-based CR programs saw significant improvements in exercise capacity and health-related quality of life scores, as well as reduced risk of hospitalization and death. Further, many studies have found no significant differences in outcomes between individuals who took part in center-based CR and those who participated in virtual/home-based CR (Molloy, et al., 2024; Krishnamurthi, et al., 2023; Tegegne, et al., 2022; Cavalheiro, et al., 2021; Piotrowicz et al., 2020; Snoek, et al., 2020; Song, et al., 2020; Jin, et al., 2019; Dorje, et al., 2019; Hwang, et al., 2017; Smolis-Bąk, et al., 2015; Lear, et al., 2014; Reid, et al., 2012).

A Cochrane systematic review and meta-analysis of 23 randomized controlled trials (n=2,890 participants) by Anderson et al. (2017b) compared the impact of home-based and center-based cardiac rehabilitation (CR) on mortality and morbidity, exercise capacity, and other outcomes in patients with heart disease. The home-based CR programs included in the review were structured, included exercise training, had clear objectives, and included monitoring, follow up visits, letters or telephone calls from staff, or at least self-monitoring diaries. The length, intensity, and specific nature of the exercise programs and monitoring varied among the programs. The control groups were center-based CR in a variety of settings (e.g. hospital physiotherapy department, university gymnasium, community sports center). Some programs (both home and control) were exercise-only, while others were comprehensive in nature. Patient inclusion criteria were age over 18 years, and one of the following: post myocardial infarction (MI), prior revascularization, and diagnosis of angina or heart failure (HF). Exclusion criteria were: patients who had undergone heart transplants, had implantable cardioverter defibrillators (ICDs) or cardiac resynchronization therapy, or who had previously undergone CR. Outcome measures varied, and included total mortality, cardiac events, exercise capacity assessed by validated outcome measure (e.g.  $VO_2$  peak, 6 minute walk test), validated measures of health-related quality of life, adherence, modifiable coronary risk factors, and costs and health service use. Follow ups typically ranged from 2-12 months, with three studies reporting data beyond 12 months. Loss to follow-up varied considerably among studies and was asymmetric across home- and center-based CR groups. Only a few trials examined the impact of losses to follow-up. Nine studies reported less than 20% attrition, four studies reported greater than 20% attrition, seven studies provided incomplete data, and three studies provided no data on attrition. Between-group outcomes up to 12 months included the following:

- Total mortality: No significant difference (based on data from 11 studies)
- Cardiac events: Not poolable, small number of studies reported data
- Exercise capacity: No significant difference
- Health-related Quality of life: Not poolable, wide variation
- Withdrawal: No significant difference (inconsistent reporting)
- Modifiable coronary risk factors: No significant difference
- Adherence: Not poolable, wide variation

- Cost and health service use: Not poolable; (home-based CR was less expensive in four studies, more expensive in one study)

Author-noted limitations included: inconsistent reporting of outcomes, considerable statistical heterogeneity across a number of outcomes among trials, short duration of most studies, and often poorly reported details of interventions making it difficult to assess whether the CR programs used would meet current standards of good practice. Additionally, there was significant heterogeneity of home CR programs as they varied in design, duration, frequency, and technological involvement (if any). Most studies included only lower-risk individuals. There was consistent underrepresentation of women. Finally, studies older than 10 years were included in the analysis.

An update to the above Cochrane systematic review and meta-analysis by Anderson et al. was published in 2023. McDonagh et al. included three new trials; a total of 24 trials (n=3046 participants) were included in the analysis. The updated review continued to find no significant differences between home- and center-based CR in the primary outcomes up to 12 months of follow-up: total mortality (risk ratio [RR] = 1.19, 95% confidence interval [CI] 0.65 to 2.16; low-certainty evidence); and exercise capacity (standardized mean difference (SMD) = -0.10, 95% CI -0.24 to 0.04; low-certainty evidence). Most studies showed no significant difference between home- and center-based CR in health-related quality of life up to 24 months follow-up. The authors concluded that both home and center-based CR, when formally supported by healthcare staff, were similarly effective in improving clinical and health-related quality of life outcomes in individuals post-myocardial infarction, post-revascularization, or with heart failure. Additional research is needed to whether the positive short-term effects of home CR programs can be confirmed in the long term (McDonagh, et al., 2023).

**Professional Societies/Organizations:** In 2019, the American Association of Cardiovascular and Pulmonary Rehabilitation (AACVPR), American Heart Association (AHA), and American College of Cardiology (ACC) issued a scientific statement on home-based cardiac rehabilitation focusing on the problem of underutilization of CR programs, despite its well-established benefits. The statement suggests one potential approach to resolving this issue is the expansion of home- or community-based CR programs, to help overcome geographic, logistical, and other access-related barriers facing traditional center-based CR programs. In particular, the suggestion is that home-based CR is a suitable and safe alternative option for CR services for stable, low- to moderate-risk patients with cardiovascular disease who lack available center-based services. Longer-term studies on the impact of home-based CR programs, as well as safety data (particularly for high-risk groups), remain lacking. Therefore thorough clinical evaluation with risk stratification is critical to ensure appropriate referral to home-based CR (Thomas, et al., 2019).

The 2024 ACC Expert Consensus Decision Pathway for the treatment of heart failure with reduced ejection fraction (HFrEF) supported the feasibility of remote telerehabilitation for patients with HFrEF who lack access to outpatient cardiac rehabilitation, citing studies that reported high adherence rates with such programs (Maddox, et al., 2024).

### **Outpatient Intensive Cardiac Rehabilitation Programs**

Several outpatient intensive cardiac rehabilitation (ICR) programs have been developed including, but not limited to, the Pritikin Program, the Ornish Program for Reversing Heart Disease, and the Benson-Henry Institute Cardiac Wellness Program (Hayes, 2018, Updated 2020; CMS, 2010; 2014). ICR are comprehensive, long-term programs involving medical evaluation, exercise, cardiac risk factor modification, education, and counseling for patients with chronic or post-acute cardiovascular disease. The Pritikin, Ornish, and Benson-Henry Institute programs are commercial, licensed products with varying program design. Common features include specific diet prescription, group support meetings, and a focus on lifestyle modification. Ongoing wellness education classes, self-guided practice, and group sessions can continue for a year or more. ICR

programs include phase III and IV elements which are considered educational and training in nature. There is a lack of large randomized prospective comparative studies in the peer-reviewed published literature that outpatient intensive cardiac rehabilitation programs improve health outcomes compared to a program of traditional outpatient cardiac rehabilitation.

A 2018 Hayes Comparative Effectiveness Review (updated 2020) evaluated the comparative effectiveness and safety of intensive cardiac rehabilitation (ICR) programs relative to usual care (UC) and conventional cardiac rehabilitation (CCR) in patients with coronary artery disease. The evidence evaluation concluded that "there is limited and very-low quality evidence, which suggests some advantages of ICR over usual care but insufficient evidence to determine whether ICR has advantages compared with conventional cardiac rehabilitation. Most evidence is based on Ornish programs, and there is an insufficient quantity of data to inform which ICR program, if any, is associated with the best outcomes."

## Medicare Coverage Determinations

	<b>Contractor</b>	<b>Determination Name/Number</b>	<b>Revision Effective Date</b>
NCD	National	Cardiac Rehabilitation Programs for Chronic Heart Failure (20.10.1)	2/18/2014
NCD	National	Intensive Cardiac Rehabilitation (ICR) Programs (20.31)	8/12/2010
NCD	National	Benson-Henry Institute Cardiac Wellness Program (20.31.3)	5/6/2014
NCD	National	The Pritikin Program (20.31.1)	8/12/2010
NCD	National	Ornish Program for Reversing Heart Disease (20.31.2)	8/12/2010
LCD		No Determination found	

Note: Please review the current Medicare Policy for the most up-to-date information.  
(NCD = National Coverage Determination; LCD = Local Coverage Determination)

## Coding Information

### Notes:

1. This list of codes may not be all-inclusive since the American Medical Association (AMA) and Centers for Medicare & Medicaid Services (CMS) code updates may occur more frequently than policy updates.
2. Deleted codes and codes which are not effective at the time the service is rendered may not be eligible for reimbursement.

**Considered Medically Necessary when criteria in the applicable policy statements listed above are met:**

<b>CPT®*</b> <b>Codes</b>	<b>Description</b>
93797	Physician or other qualified health care professional services for outpatient cardiac rehabilitation; without continuous ECG monitoring (per session)
93798	Physician or other qualified health care professional services for outpatient cardiac rehabilitation; with continuous ECG monitoring (per session)

**Considered Not Medically Necessary:**

<b>HCPCS Codes</b>	<b>Description</b>
G0422	Intensive cardiac rehabilitation; with or without continuous ECG monitoring with exercise, per session
G0423	Intensive cardiac rehabilitation; with or without continuous ECG monitoring, without exercise, per session
S9472	Cardiac rehabilitation program, non-physician provider, per diem

**\*Current Procedural Terminology (CPT®) ©2024 American Medical Association: Chicago, IL.**

## References

1. Abraham LN, Sibilitz KL, Berg SK, Tang LH, Risom SS, Lindschou J, Taylor RS, Borregaard B, Zwisler AD. Exercise-based cardiac rehabilitation for adults after heart valve surgery. *Cochrane Database Syst Rev.* 2021 May 7;5(5):CD010876.
2. Anderson L, Nguyen TT, Dall CH, Burgess L, Bridges C, Taylor RS. Exercise-based cardiac rehabilitation in heart transplant recipients. *Cochrane Database Syst Rev.* 2017a Apr 4;4(4):CD012264.
3. Anderson L, Sharp GA, Norton RJ, Dalal H, Dean SG, Jolly K, Cowie A, Zawada A, Taylor RS. Home-based versus centre-based cardiac rehabilitation. *Cochrane Database of Systematic Reviews* 2017b, Issue 6. Art. No.: CD007130.
4. Anderson L, Taylor RS. Cardiac rehabilitation for people with heart disease: an overview of Cochrane systematic reviews. *Cochrane Database Syst Rev.* 2014 Dec 12;2014(12):CD011273.
5. Balady GJ, Williams MA, Ades PA, Bittner V, Comoss P, Foody JM, Franklin B, Sanderson B, Southard, D. Core components of cardiac rehabilitation/secondary prevention programs: 2007 update: a scientific statement from the American Heart Association Exercise, Cardiac Rehabilitation, and Prevention Committee, the Council on Clinical Cardiology; the Councils on Cardiovascular Nursing, Epidemiology and Prevention, and Nutrition, Physical Activity, and Metabolism; and the American Association of Cardiovascular and Pulmonary Rehabilitation. *Circulation.* 2007 May 22;115(20):2675-82.
6. Bartels MN, Lehman YJ, Rashbaum IG. Cardiac rehabilitation: Lifestyle, exercise & risk factors modification. In: O' Young BJ, Young MA, Stiens SA, Wu SSH. *Physical Medicine and Rehabilitation Secrets.* Philadelphia, PA: Elsevier; 2024. 445-451.
7. Beatty AL, Beckie TM, Dodson J, Goldstein CM, Hughes JW, Kraus WE, Martin SS, Olson TP, Pack QR, Stolp H, Thomas RJ, Wu WC, Franklin BA. A New Era in Cardiac Rehabilitation Delivery: Research Gaps, Questions, Strategies, and Priorities. *Circulation.* 2023 Jan 17;147(3):254-266. Epub 2023 Jan 17. Erratum in: *Circulation.* 2023 Feb 28;147(9):e648.
8. Brown TM, Pack QR, Aberegg E, Brewer LC, Ford YR, Forman DE, Gathright EC, Khadanga S, Ozemek C, Thomas RJ; American Heart Association Exercise, Cardiac Rehabilitation and Secondary Prevention Committee of the Council on Clinical Cardiology; Council on Cardiovascular and Stroke Nursing; Council on Lifestyle and Cardiometabolic Health; and Council on Quality of Care and Outcomes Research. Core Components of Cardiac

Rehabilitation Programs: 2024 Update: A Scientific Statement From the American Heart Association and the American Association of Cardiovascular and Pulmonary Rehabilitation. *Circulation*. 2024 Oct 29;150(18):e328-e347.

9. Buckley BJ, Long L, Risom SS, Lane DA, Berg SK, Gluud C, Palm P, Sibilitz KL, Svendsen JH, Zwisler AD, Lip GY, Neubeck L, Taylor RS. Exercise-based cardiac rehabilitation for adults with atrial fibrillation. *Cochrane Database Syst Rev*. 2024 Sep 17;9(9):CD011197
10. Cai C, Bao Z, Wu N, Wu F, Sun G, Yang G, Chen M. A novel model of home-based, patient-tailored and mobile application-guided cardiac telerehabilitation in patients with atrial fibrillation: A randomised controlled trial. *Clin Rehabil*. 2022 Jan;36(1):40-50.
11. Cavalheiro AH, Silva Cardoso J, Rocha A, Moreira E, Azevedo LF. Effectiveness of Tele-rehabilitation Programs in Heart Failure: A Systematic Review and Meta-analysis. *Health Serv Insights*. 2021;14:11786329211021668. Published 2021 Jun 15.
12. Centers for Medicare and Medicaid Services (CMS). Decision Memo for Cardiac Rehabilitation Programs (CAG-00089R2). Feb 22, 2010. Accessed Mar 25, 2025. Available at URL address: <https://www.cms.gov/medicare-coverage-database/view/ncacal-decision-memo.aspx?proposed=N&NCAId=241>
13. Centers for Medicare and Medicaid Services (CMS). Decision Memo for Cardiac Rehabilitation (CR) Programs - Chronic Heart Failure (CAG-00437N). Feb 18, 2014. Accessed Mar 25, 2025. Available at URL address: <https://www.cms.gov/medicare-coverage-database/view/ncacal-decision-memo.aspx?proposed=N&NCAId=270>
14. Centers for Medicare and Medicaid Services (CMS). Medicare Claims Processing Manual. Ch. 32 Billing Requirements for Special Services. Issued Oct 11, 2024. Accessed Mar 25, 2025. Available at URL address: <https://www.cms.gov/regulations-and-guidance/guidance/manuals/downloads/clm104c32.pdf>
15. Centers for Medicare and Medicaid Services (CMS). Medicare Coverage Database. Accessed Mar 25, 2025. Available at URL address: <https://www.cms.gov/medicare-coverage-database/search.aspx>
16. Centers for Medicare and Medicaid Services (CMS). National Coverage Determination (NCD) for Benson-Henry Institute Cardiac Wellness Program (20.31.3). Accessed Mar 25, 2025. Available at URL address: <https://www.cms.gov/medicare-coverage-database/view/ncd.aspx?ncdid=362>
17. Centers for Medicare and Medicaid Services (CMS). National Coverage Determination (NCD) for Cardiac Rehabilitation Programs for Chronic Heart Failure (20.10.1). Accessed Mar 25, 2025. Available at URL address: <https://www.cms.gov/medicare-coverage-database/view/ncd.aspx?ncdid=359>
18. Centers for Medicare and Medicaid Services (CMS). National Coverage Determination (NCD) for Intensive Cardiac Rehabilitation (ICR) Programs (20.31). Accessed Mar 25, 2025. Available at URL address: <https://www.cms.gov/medicare-coverage-database/view/ncd.aspx?ncdid=339>
19. Centers for Medicare and Medicaid Services (CMS). National Coverage Determination (NCD) for Ornish Program for Reversing Heart Disease (20.31.2). Accessed Mar 25, 2025.

Available at URL address: <https://www.cms.gov/medicare-coverage-database/view/ncd.aspx?ncdid=341>

20. Centers for Medicare and Medicaid Services (CMS). National Coverage Determination (NCD) for The Pritikin Program (20.31.1). Accessed Mar 25, 2025. Available at URL address: <https://www.cms.gov/medicare-coverage-database/view/ncd.aspx?ncdid=340>
21. Clark AM, Hartling L, Vandermeer B, McAlister FA. Meta-analysis: secondary prevention programs for patients with coronary artery disease. *Ann Intern Med.* 2005 Nov 1;143(9):659–72.
22. Cruz-Cobo C, Bernal-Jiménez MÁ, Vázquez-García R, Santi-Cano MJ. Effectiveness of mHealth Interventions in the Control of Lifestyle and Cardiovascular Risk Factors in Patients After a Coronary Event: Systematic Review and Meta-analysis. *JMIR Mhealth Uhealth.* 2022 Dec 2;10(12):e39593.
23. Davis AM. Cardiac Rehabilitation. In: Frontera WR, Silver JK, Rizzo TD. *Essentials of Physical Medicine and Rehabilitation.* 4<sup>th</sup> ed. Philadelphia PA: Elsevier. 2019. Ch 123, 678-683.
24. Dibben G, Faulkner J, Oldridge N, Rees K, Thompson DR, Zwisler AD, Taylor RS. Exercise-based cardiac rehabilitation for coronary heart disease. *Cochrane Database Syst Rev.* 2021 Nov 6;11(11):CD001800.
25. Dorje T, Zhao G, Tso K, Wang J, Chen Y, Tsokey L, Tan BK, Scheer A, Jacques A, Li Z, Wang R, Chow CK, Ge J, Maiorana A. Smartphone and social media-based cardiac rehabilitation and secondary prevention in China (SMART-CR/SP): a parallel-group, single-blind, randomised controlled trial. *Lancet Digit Health.* 2019 Nov;1(7):e363-e374.
26. Ghisi GLM, Kin SMR, Price J, Beckie TM, Mamataz T, Naheed A, Grace SL. Women-Focused Cardiovascular Rehabilitation: An International Council of Cardiovascular Prevention and Rehabilitation Clinical Practice Guideline. *Can J Cardiol.* 2022 Dec;38(12):1786-1798.
27. Golbus JR, Lopez-Jimenez F, Barac A, Cornwell WK 3rd, Dunn P, Forman DE, Martin SS, Schorr EN, Supervia M; Exercise, Cardiac Rehabilitation and Secondary Prevention Committee of the Council on Clinical Cardiology; Council on Lifelong Congenital Heart Disease and Heart Health in the Young; Council on Quality of Care and Outcomes Research; and Council on Cardiovascular and Stroke Nursing. Digital Technologies in Cardiac Rehabilitation: A Science Advisory From the American Heart Association. *Circulation.* 2023 Jul 4;148(1):95-107.
28. Gore S, Khanna H, Kumar A. Effects of Comprehensive Outpatient Cardiac Rehabilitation on Exercise Capacity, Functional Status, and Quality of Life in People With Heart Failure: A Systematic Review and Meta-Analysis. *Phys Ther.* 2023 Nov 4;103(11):pzad119.
29. Halperin JL, Levine GN, Al-Khatib SM, Birtcher KK, Bozkurt B, Brindis RG, Cigarroa JE, Curtis LH, Fleisher LA, Gentile F, Gidding S, Hlatky MA, Ikonomidis J, Joglar J, Pressler SJ, Wijeyesundera DN. Further Evolution of the ACC/AHA Clinical Practice Guideline Recommendation Classification System: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol.* 2016 Apr 5;67(13):1572-1574.

30. Hamm LF, Sanderson BK, Ades PA, Berra K, Kaminsky LA, Roitman JL, Williams MA. Core competencies for cardiac rehabilitation/secondary prevention professionals: 2010 update: position statement of the American Association of Cardiovascular and Pulmonary Rehabilitation. *J Cardiopulm Rehabil Prev.* 2011 Jan-Feb;31(1):2-10.
31. Hammill BG, Curtis LH, Schulman KA, Whellan DJ. Relationship between cardiac rehabilitation and long-term risks of death and myocardial infarction among elderly Medicare beneficiaries. *Circulation.* 2010 Jan 5;121(1):63-70.
32. Hayes, Inc. Medical Technology Directory. Comparative Effectiveness Review. Intensive Cardiac Rehabilitation Programs for Coronary Artery Disease. Hayes, Inc.; published Feb 22, 2018, Updated May 27, 2020.
33. Heidenreich PA, Bozkurt B, Aguilar D, Allen LA, Byun JJ, Colvin MM, Deswal A, Drazner MH, Dunlay SM, Evers LR, Fang JC, Fedson SE, Fonarow GC, Hayek SS, Hernandez AF, Khazanie P, Kittleson MM, Lee CS, Link MS, Milano CA, Nwacheta LC, Sandhu AT, Stevenson LW, Vardeny O, Vest AR, Yancy CW. 2022 AHA/ACC/HFSA guideline for the management of heart failure: a report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation.* 2022;145:e895–e1032.
34. Husaini M, Deych E, Waken RJ, Sells B, Lai A, Racette SB, Rich MW, Joynt Maddox KE, Peterson LR. Intensive Versus Traditional Cardiac Rehabilitation: Mortality and Cardiovascular Outcomes in a 2016-2020 Retrospective Medicare Cohort. *Circ Cardiovasc Qual Outcomes.* 2023 Dec;16(12):e010131.
35. Hwang R, Bruning J, Morris NR, Mandrusiak A, Russell T. Home-based telerehabilitation is not inferior to a centre-based program in patients with chronic heart failure: a randomised trial. *Journal of Physiotherapy.* 2017 63: 101–107
36. Jepma P, Verweij L, Buurman BM, Terbraak MS, Daliri S, Latour CHM, Ter Riet G, Karapinar-Çarkit F, Dekker J, Klunder JL, Liem SS, Moons AHM, Peters RJG, Scholte Op Reimer WJM. The nurse-coordinated cardiac care bridge transitional care programme: a randomised clinical trial. *Age Ageing.* 2021 Nov 10;50(6):2105-2115.
37. Ji H, Fang L, Yuan L, Zhang Q. Effects of Exercise-Based Cardiac Rehabilitation in Patients With Acute Coronary Syndrome: A Meta-Analysis. *Meta-Analysis Med Sci Monit.* 25, 5015-5027 2019 Jul 7
38. Jin K, Khonsari S, Gallagher R, Gallagher P, Clark AM, Freedman B, Briffa T, Bauman A, Redfern J, Neubeck L. Telehealth interventions for the secondary prevention of coronary heart disease: A systematic review and meta-analysis. *Eur J Cardiovasc Nurs.* 2019 Apr;18(4):260-271.
39. Jolly K, Taylor R, Lip GY, Greenfield S, Raftery J, Mant J, Lane D, Jones M, Lee KW, Stevens A. The Birmingham Rehabilitation Uptake Maximisation Study (BRUM). Home-based compared with hospital-based cardiac rehabilitation in a multi-ethnic population: cost-effectiveness and patient adherence. *Health Technol Assess.* 2007 Sep;11(35):1-118.
40. Krishnamurthi N, Schopfer DW, Shen H, Rohrbach G, Elnaggar A, Whooley MA. Association of Home-Based Cardiac Rehabilitation With Lower Mortality in Patients With Cardiovascular Disease: Results From the Veterans Health Administration Healthy Heart Program. *J Am Heart Assoc.* 2023 Mar 7;12(5):e025856.

41. La Rovere MT, Traversi E. Role and Efficacy of Cardiac Rehabilitation in Patients With Heart Failure. *Monaldi Arch Chest Dis*, 89 (1) 2019 Apr 12
42. Lakhani F, Racette SB, Park LK, Deych E, Williams D, McKenzie KM, Stranczek NA, McKenzie EJ, Lenze EJ, Durbin D, Jonagan J, Carson T, Talpade N, Rich MW, Fuentes LL, Peterson LR. Prospective Study of the Impact of Outpatient Intensive Cardiac Rehabilitation on Diet Quality, Health-related Quality of Life, and Cardiovascular Health Indices. *Am J Cardiol*. 2023 Apr 1;192:60-66.
43. Lawler PR, Filion KB, Eisenberg MJ. Efficacy of exercise-based cardiac rehabilitation post-myocardial infarction: a systematic review and meta-analysis of randomized controlled trials. *Am Heart J*. 2011 Oct;162(4):571-584.e2.
44. Lawton JS, Tamis-Holland JE, Bangalore S, Bates ER, Beckie TM, Bischoff JM, Bittl JA, Cohen MG, DiMaio JM, Don CW, Fremes SE, Gaudino MF, Goldberger ZD, Grant MC, Jaswal JB, Kurlansky PA, Mehran R, Metkus TS Jr, Nnacheta LC, Rao SV, Sellke FW, Sharma G, Yong CM, Zwischenberger BA. 2021 ACC/AHA/SCAI Guideline for Coronary Artery Revascularization: Executive Summary: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation*. 2022 Jan 18;145(3):e4-e17. Erratum in: *Circulation*. 2022 Mar 15;145(11):e771.
45. Lear SA, Singer J, Banner-Lukaris D, Horvat D, Park JE, Bates J, Ignaszewski A. Randomized trial of a virtual cardiac rehabilitation program delivered at a distance via the Internet. *Circ Cardiovasc Qual Outcomes*. 2014 Nov;7(6):952-9.
46. Leon AS, Franklin BA, Costa F, Balady GJ, Berra KA, Stewart KJ, Thompson PD, Williams MA, Lauer MS. Cardiac rehabilitation and secondary prevention of coronary heart disease: an American Heart Association scientific statement from the Council on Clinical Cardiology (Subcommittee on Exercise, Cardiac Rehabilitation, and Prevention) and the Council on Nutrition, Physical Activity, and Metabolism (Subcommittee on Physical Activity), in collaboration with the American association of Cardiovascular and Pulmonary Rehabilitation. *Circulation*. 2005 Jan 25;111(3):369-76.
47. Li Z, Hui Z, Zheng Y, Yu J, Zhang J. Efficacy of Phase II Remote Home Rehabilitation in Patients with Acute Myocardial Infarction after Percutaneous Coronary Intervention. *Contrast Media Mol Imaging*. 2022 Jun 13;2022:4634769.
48. Long L, Anderson L, Dewhirst AM, He J, Bridges C, Gandhi M, Taylor RS. Exercise-based cardiac rehabilitation for adults with stable angina. *Cochrane Database Syst Rev*. 2018 Feb 2;2(2):CD012786.
49. Maddox TM, Januzzi JL Jr, Allen LA, Breathett K, Brouse S, Butler J, Davis LL, Fonarow GC, Ibrahim NE, Lindenfeld J, Masoudi FA, Motiwala SR, Oliveros E, Walsh MN, Wasserman A, Yancy CW, Youmans QR. 2024 ACC Expert Consensus Decision Pathway for Treatment of Heart Failure With Reduced Ejection Fraction: A Report of the American College of Cardiology Solution Set Oversight Committee. *J Am Coll Cardiol*. 2024 Apr 16;83(15):1444-1488.
50. Mathews L, Brewer LC. A Review of Disparities in Cardiac Rehabilitation: Evidence, Drivers, and Solutions. *J Cardiopulm Rehabil Prev*. 2021 Nov 1;41(6):375-382.

51. McDonagh ST, Dalal H, Moore S, Clark CE, Dean SG, Jolly K, Cowie A, Afzal J, Taylor RS. Home-based versus centre-based cardiac rehabilitation. *Cochrane Database Syst Rev*. 2023 Oct 27;10(10):CD007130.
52. Molloy CD, Long L, Mordi IR, Bridges C, Sagar VA, Davies EJ, Coats AJS, Dalal H, Rees K, Singh SJ, Taylor RS. Exercise-based cardiac rehabilitation for adults with heart failure - 2023 Cochrane systematic review and meta-analysis. *Eur J Heart Fail*. 2023 Dec;25(12):2263-2273.
53. Mosca L, Benjamin EJ, Berra K, Bezanson JL, Dolor RJ, Lloyd-Jones DM, Newby LK, Piña IL, Roger VL, Shaw LJ, Zhao D, Beckie TM, Bushnell C, D'Armiento J, Kris-Etherton PM, Fang J, Ganiats TG, Gomes AS, Gracia CR, Haan CK, Jackson EA, Judelson DR, Kelepouris E, Lavie CJ, Moore A, Nussmeier NA, Ofili E, Oparil S, Ouyang P, Pinn VW, Sherif K, Smith SC Jr, Sopko G, Chandra-Strobos N, Urbina EM, Vaccarino V, Wenger NK. Effectiveness-based guidelines for the prevention of cardiovascular disease in women--2011 update: a guideline from the American Heart Association. *Circulation*. 2011 Mar 22;123(11):1243-62. Erratum in: *Circulation*. 2011 Jun 7;123(22):e624. Erratum in: *Circulation*. 2011 Oct 18;124(16):e427.
54. Nielsen KM, Zwisler AD, Taylor RS, Svendsen JH, Lindschou J, Anderson L, Jakobsen JC, Berg SK. Exercise-based cardiac rehabilitation for adult patients with an implantable cardioverter defibrillator. *Cochrane Database Syst Rev*. 2019 Feb 12;2(2):CD011828.
55. Oehler A, Battel L, Poornima IG. FACCE Expert Analysis. Cardiac rehabilitation and health care disparities in the post-COVID-19 era. Sep 25, 2023. Accessed Mar 27, 2025. Available at URL address: <https://www.acc.org/Latest-in-Cardiology/Articles/2023/09/25/14/05/Cardiac-Rehabilitation-and-Health-Care-Disparities-in-the-Post-COVID-19-Era>
56. O'Gara PT, Kushner FG, Ascheim DD, Casey DE Jr, Chung MK, de Lemos JA, et al 2013 ACCF/AHA Guideline for the Management of ST-Elevation Myocardial Infarction: A Report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. 2013 Jan 29;61(4):e78-e140.
57. Orzechowski P, Piotrowicz R, Zaręba W, Głowczyńska R, Szalewska D, Pluta S, Irzmański R, Kalarus Z, Banach M, Opolski G, Pencina MJ, Kowalik I, Piotrowicz E. Assessment of ECG during hybrid comprehensive telerehabilitation in heart failure patients-Subanalysis of the Telerehabilitation in Heart Failure Patients (TELEREH-HF) randomized clinical trial. *Ann Noninvasive Electrocardiol*. 2021 Nov;26(6):e12887.
58. Patel P, Dhindsa D, Eapen DJ, Khera A, Gulati M, Stone NJ, Yancy CW, Rumsfeld JS, Sperling LS. Optimizing the Potential for Telehealth in Cardiovascular Care (in the Era of COVID-19): Time Will Tell. *Am J Med*. 2021 Aug;134(8):945-951.
59. Pio CSA, Chaves G, Davies P, Taylor R, Grace S. Interventions to Promote Patient Utilization of Cardiac Rehabilitation: Cochrane Systematic Review and Meta-Analysis. *J Clin Med*. 2019 Feb 5;8(2):189.
60. Piotrowicz E, Pencina MJ, Opolski G, et al. Effects of a 9-Week Hybrid Comprehensive Telerehabilitation Program on Long-term Outcomes in Patients With Heart Failure: The Telerehabilitation in Heart Failure Patients (TELEREH-HF) Randomized Clinical Trial. *JAMA Cardiol*. 2020;5(3):300-308.

61. Purcell C, Dibben G, Hilton Boon M, Matthews L, Palmer VJ, Thomson M, Smillie S, Simpson SA, Taylor RS. Social network interventions to support cardiac rehabilitation and secondary prevention in the management of people with heart disease. *Cochrane Database Syst Rev*. 2023 Jun 28;6(6):CD013820.
62. Rao SV, O'Donoghue ML, Ruel M, Rab T, Tamis-Holland JE, Alexander JH, Baber U, Baker H, Cohen MG, Cruz-Ruiz M, Davis LL, de Lemos JA, DeWald TA, Elgendy IY, Feldman DN, Goyal A, Isiadinso I, Menon V, Morrow DA, Mukherjee D, Platz E, Promes SB, Sandner S, Sandoval Y, Schunder R, Shah B, Stopyra JP, Talbot AW, Taub PR, Williams MS. 2025 ACC/AHA/ACEP/NAEMSP/SCAI Guideline for the Management of Patients With Acute Coronary Syndromes: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation*. 2025 Feb 27.
63. Ramachandran HJ, Jiang Y, Tam WWS, Yeo TJ, Wang W. Effectiveness of home-based cardiac telerehabilitation as an alternative to Phase 2 cardiac rehabilitation of coronary heart disease: a systematic review and meta-analysis. *Eur J Prev Cardiol*. 2021 Jul 13:zwab106.
64. Rawstorn JC, Gant N, Direito A, Beckmann C, Maddison R. Telehealth exercise-based cardiac rehabilitation: a systematic review and meta-analysis. *Heart*. 2016 Aug 1;102(15):1183-92.
65. Reid RD, Morrin LI, Beaton LJ, Papadakis S, Kocourek J, McDonnell L, Slovinec D'Angelo ME, Tulloch H, Suskin N, Unsworth K, Blanchard C, Pipe AL. Randomized trial of an internet-based computer-tailored expert system for physical activity in patients with heart disease. *Eur J Prev Cardiol*. 2012 Dec;19(6):1357-64.
66. Sachdev V, Sharma K, Keteyian SJ, Alcaín CF, Desvigne-Nickens P, Fleg JL, Florea VG, Franklin BA, Guglin M, Halle M, Leifer ES, Panjrath G, Tinsley EA, Wong RP, Kitzman DW; American Heart Association Heart Failure and Transplantation Committee of the Council on Clinical Cardiology; Council on Arteriosclerosis, Thrombosis and Vascular Biology; and American College of Cardiology. Supervised Exercise Training for Chronic Heart Failure With Preserved Ejection Fraction: A Scientific Statement From the American Heart Association and American College of Cardiology. *Circulation*. 2023 Mar 21.
67. Sawyer KN, Camp-Rogers TR, Kotini-Shah P, Del Rios M, Gossip MR, Moitra VK, Haywood KL, Dougherty CM, Lubitz SA, Rabinstein AA, Rittenberger JC, Callaway CW, Abella BS, Geocadin RG, Kurz MC; American Heart Association Emergency Cardiovascular Care Committee; Council on Cardiovascular and Stroke Nursing; Council on Genomic and Precision Medicine; Council on Quality of Care and Outcomes Research; and Stroke Council. Sudden Cardiac Arrest Survivorship: A Scientific Statement From the American Heart Association. *Circulation*. 2020 Mar 24;141(12):e654-e685.
68. Smith KM, McKelvie RS, Thorpe KE, Arthur HM. Six-year follow-up of a randomised controlled trial examining hospital versus home-based exercise training after coronary artery bypass graft surgery. *Heart*. 2011 Jul;97(14):1169-74.
69. Smith SC Jr, Benjamin EJ, Bonow RO, Braun LT, Creager MA, Franklin BA, Gibbons RJ, Grundy SM, Hiratzka LF, Jones DW, Lloyd-Jones DM, Minissian M, Mosca L, Peterson ED, Sacco RL, Spertus J, Stein JH, Taubert KA; World Heart Federation and the Preventive Cardiovascular Nurses Association. AHA/ACCF Secondary Prevention and Risk Reduction Therapy for Patients with Coronary and other Atherosclerotic Vascular Disease: 2011 update: a guideline from the American Heart Association and American College of

Cardiology Foundation. *Circulation*. 2011 Nov 29;124(22):2458-73. Erratum in: *Circulation*. 2015 Apr 14;131(15):e408.

70. Smolis-Bąk E, Dąbrowski R, Piotrowicz E, Chwyczko T, Dobraszkievicz-Wasilewska B, Kowalik I, Kazimierska B, Jędrzejczyk B, Smolis R, Gepner K, Maciąg A, Sterliński M, Szwed H. Hospital-based and telemonitoring guided home-based training programs: effects on exercise tolerance and quality of life in patients with heart failure (NYHA class III) and cardiac resynchronization therapy. A randomized, prospective observation. *Int J Cardiol*. 2015 Nov 15;199:442-7.
71. Snoek JA, Prescott EI, van der Velde AE, Eijsvogels TMH, Mikkelsen N, Prins LF, Bruins W, Meindersma E, González-Juanatey JR, Peña-Gil C, González-Salvado V, Moatemri F, Iliou MC, Marcin T, Eser P, Wilhelm M, Van't Hof AWJ, de Kluiver EP. Effectiveness of Home-Based Mobile Guided Cardiac Rehabilitation as Alternative Strategy for Nonparticipation in Clinic-Based Cardiac Rehabilitation Among Elderly Patients in Europe: A Randomized Clinical Trial. *JAMA Cardiol*. 2020 Oct 28:e205218
72. Song Y, Ren C, Liu P, Tao L, Zhao W, Gao W. Effect of Smartphone-Based Telemonitored Exercise Rehabilitation among Patients with Coronary Heart Disease. *J Cardiovasc Transl Res*. 2020;13(4):659-667.
73. Szalewska D, Głowczyńska R, Piotrowicz R, Kowalik I, Pencina MJ, Opolski G, Zaręba W, Banach M, Orzechowski P, Pluta S, Irzmański R, Kalarus Z, Piotrowicz E. An aetiology-based subanalysis of the Telerehabilitation in Heart Failure Patients (TELEREH-HF) trial. *ESC Heart Fail*. 2021 Apr;8(2):1263-1273.
74. Taylor RS, Brown A, Ebrahim S, Jolliffe J, Noorani H, Rees K, et al. Exercise-based rehabilitation for patients with coronary heart disease: systematic review and meta-analysis of randomized controlled trials. *Am J Med*. 2004 May 15;116(10):682-92.
75. Tegegne TK, Rawstorn JC, Nourse RA, Kibret KT, Ahmed KY, Maddison R. Effects of exercise-based cardiac rehabilitation delivery modes on exercise capacity and health-related quality of life in heart failure: a systematic review and network meta-analysis. *Open Heart*. 2022 Jun;9(1):e001949.
76. Thomas RJ. Comprehensive Cardiac Rehabilitation. In: Libby P, Bonow RO, Mann DL, Tomaselli GF, Bhatt DL. *Braunwald's Heart Disease: A Textbook of Cardiovascular Medicine*. 12th ed. Philadelphia, PA: Elsevier; 2022. 588-592.
77. Thomas RJ, Beatty AL, Beckie TM, Brewer LC, Brown TM, Forman DE, Franklin BA, Keteyian SJ, Kitzman DW, Regensteiner JG, Sanderson BK; Whooley MA. Home-based cardiac rehabilitation: a scientific statement from the American Association of Cardiovascular and Pulmonary Rehabilitation, the American Heart Association, and the American College of Cardiology. *Circulation*. 2019;140:e69–e89.
78. Thomas RJ, King M, Lui K, Oldridge N, Piña IL, Spertus J. AACVPR/ACC/AHA 2007 performance measures on cardiac rehabilitation for referral to and delivery of cardiac rehabilitation/secondary prevention services. *Circulation*. 2007;116:1611–1642
79. Thomas RJ, King M, Lui K, Oldridge N, Piña IL, Spertus J. AACVPR/ACCF/AHA 2010 update: performance measures on cardiac rehabilitation for referral to cardiac rehabilitation/secondary prevention services: a report of the American Association of Cardiovascular and Pulmonary Rehabilitation and the American College of Cardiology

Foundation/American Heart Association Task Force on Performance Measures (Writing Committee to Develop Clinical Performance Measures for Cardiac Rehabilitation). *Circulation*. 2010;122:1342–1350.

80. Thomas RJ, Balady G, Banka G, Beckie TM, Chiu J, Gokak S, Ho PM, Keteyian SJ, King M, Lui K, Pack Q, Sanderson BK, Wang TY. 2018 ACC/AHA clinical performance and quality measures for cardiac rehabilitation: a report of the American College of Cardiology/American Heart Association Task Force on Performance Measures. *Circ Cardiovasc Qual Outcomes*. 2018;11:e000037.
81. Verweij L, Petri ACM, MacNeil-Vroomen JL, Jepma P, Latour CHM, Peters RJG, Scholte Op Reimer WJM, Buurman BM, Bosmans JE. The Cardiac Care Bridge transitional care program for the management of older high-risk cardiac patients: An economic evaluation alongside a randomized controlled trial. *PLoS One*. 2022 Jan 27;17(1):e0263130.
82. Virani SS, Newby LK, Arnold SV, Bittner V, Brewer LC, Demeter SH, Dixon DL, Fearon WF, Hess B, Johnson HM, Kazi DS, Kolte D, Kumbhani DJ, LoFaso J, Mahtta D, Mark DB, Minissian M, Navar AM, Patel AR, Piano MR, Rodriguez F, Talbot AW, Taqueti VR, Thomas RJ, van Diepen S, Wiggins B, Williams MS; Peer Review Committee Members. 2023 AHA/ACC/ACCP/ASPC/NLA/PCNA Guideline for the Management of Patients With Chronic Coronary Disease: A Report of the American Heart Association/American College of Cardiology Joint Committee on Clinical Practice Guidelines. *Circulation*. 2023 Aug 29;148(9):e9-e119. Erratum in: *Circulation*. 2023 Sep 26;148(13):e148. Erratum in: *Circulation*. 2023 Dec 5;148(23):e186.

## Revision Details

Type of Revision	Summary of Changes	Date
Annual Review	<ul style="list-style-type: none"> <li>Revised policy statement for cardiac rehabilitation.</li> </ul>	5/15/2025
Annual Review	<ul style="list-style-type: none"> <li>Removed “center-based” from policy statement; removed noncoverage policy statement for virtual/remote home-based and hybrid cardiac rehabilitation.</li> </ul>	6/15/2024

---

“Cigna Companies” refers to operating subsidiaries of The Cigna Group. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life Insurance Company, Connecticut General Life Insurance Company, Evernorth Behavioral Health, Inc., Cigna Health Management, Inc., and HMO or service company subsidiaries of The Cigna Group. © 2025 The Cigna Group.