



Drug Coverage Policy

Effective Date5/15/2025

Coverage Policy NumberIP0564

Policy Title Altuviio

Hemophilia – Altuviio

- Altuviio™ (antihemophilic factor [recombinant] Fc-VWF-XTEN fusion protein-ehtl intravenous infusion - Bioverativ/Sanofi)

INSTRUCTIONS FOR USE

The following Coverage Policy applies to health benefit plans administered by Cigna Companies. Certain Cigna Companies and/or lines of business only provide utilization review services to clients and do not make coverage determinations. References to standard benefit plan language and coverage determinations do not apply to those clients. Coverage Policies are intended to provide guidance in interpreting certain standard benefit plans administered by Cigna Companies. Please note, the terms of a customer's particular benefit plan document [Group Service Agreement, Evidence of Coverage, Certificate of Coverage, Summary Plan Description (SPD) or similar plan document] may differ significantly from the standard benefit plans upon which these Coverage Policies are based. For example, a customer's benefit plan document may contain a specific exclusion related to a topic addressed in a Coverage Policy. In the event of a conflict, a customer's benefit plan document always supersedes the information in the Coverage Policies. In the absence of a controlling federal or state coverage mandate, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of 1) the terms of the applicable benefit plan document in effect on the date of service; 2) any applicable laws/regulations; 3) any relevant collateral source materials including Coverage Policies and; 4) the specific facts of the particular situation. Each coverage request should be reviewed on its own merits. Medical directors are expected to exercise clinical judgment where appropriate and have discretion in making individual coverage determinations. Where coverage for care or services does not depend on specific circumstances, reimbursement will only be provided if a requested service(s) is submitted in accordance with the relevant criteria outlined in the applicable Coverage Policy, including covered diagnosis and/or procedure code(s). Reimbursement is not allowed for services when billed for conditions or diagnoses that are not covered under this Coverage Policy (see "Coding Information" below). When billing, providers must use the most appropriate codes as of the effective date of the submission. Claims submitted for services that are not accompanied by covered code(s) under the applicable Coverage Policy will be denied as not covered. Coverage Policies relate exclusively to the administration of health benefit plans. Coverage Policies are not recommendations for treatment and should never be used as treatment guidelines. In certain markets, delegated vendor guidelines may be used to support medical necessity and other coverage determinations.

OVERVIEW

Altuviio, a recombinant DNA-derived Factor VIII concentrate, is indicated for use in **hemophilia A** in adults and children for:¹

- **Routine prophylaxis** to reduce the frequency of bleeding episodes.
- **On-demand treatment and control** of bleeding episodes.
- **Perioperative management** of bleeding.

It is notable that Altuviiiio has demonstrated a 3- to 4-fold prolonged half-life relative to other standard and extended half-life products.¹

Disease Overview

Hemophilia A is an X-linked bleeding disorder primarily impacting males caused by a deficiency in Factor VIII.²⁻⁵ In the US, the incidence of hemophilia A in males is 1:5,000 with an estimated 20,000 people in the US living with hemophilia A. The condition is characterized by bleeding in joints, either spontaneously or in a provoked joint by trauma. Bleeding can occur in many different body areas as well (e.g., muscles, central nervous system). The bleeding manifestations can lead to substantial morbidity such as hemophilic arthropathy. Disease severity is usually defined by the plasma levels or activity of Factor VIII classified as follows: severe (levels < 1% of normal), moderate (levels 1% to 5% of normal), and mild (levels > 5% to < 40% of normal); phenotypic expression may vary. Approximately 50% of patients with hemophilia A are categorized as having severe disease.

Guidelines

Guidelines for hemophilia from the National Bleeding Disorders Foundation (October 2024)⁶ recognize Altuviiiio as a product with a prolonged half-life that differs from other recombinant Factor VIII concentrates.

Dosing Considerations

Dosing of clotting factor concentrates is highly individualized. The National Hemophilia Foundation's Medical and Scientific Advisory Council (MASAC) provides recommendations regarding doses of clotting factor concentrate in the home (2016).⁷ The number of required doses varies greatly and is dependent on the severity of the disorder and the prescribed regimen. Per MASAC guidance, patients on prophylaxis should also have a minimum of one major dose and two minor doses on hand for breakthrough bleeding in addition to the prophylactic doses used monthly. The guidance also notes that an adequate supply of clotting factor concentrate is needed to accommodate weekends and holidays. Therefore, maximum doses in this policy allow for prophylactic dosing plus three days of acute bleeding or perioperative management per 28 days. Doses exceeding this quantity will be reviewed on a case-by-case basis by a clinician.

Coverage Policy

Policy Statement

Prior Authorization is recommended for prescription benefit coverage of Altuviiiio. All approvals are provided for the duration noted below. In cases where the approval is authorized in months, 1 month is equal to 30 days. Because of the specialized skills required for evaluation and diagnosis of patients treated with Altuviiiio as well as the monitoring required for adverse events and long-term efficacy, approval requires Altuviiiio to be prescribed by a physician who has consulted with or who specializes in the condition.

Altuviiiio, a recombinant DNA-derived Factor VIII concentrate, is indicated for use in **hemophilia A** in adults and children for:¹

- **Routine prophylaxis** to reduce the frequency of bleeding episodes.
- **On-demand treatment and control** of bleeding episodes.
- **Perioperative management** of bleeding.

It is notable that Altuviiiio has demonstrated a 3- to 4-fold prolonged half-life relative to other standard and extended half-life products.¹

Disease Overview

Hemophilia A is an X-linked bleeding disorder primarily impacting males caused by a deficiency in Factor VIII.²⁻⁵ In the US, the incidence of hemophilia A in males is 1:5,000 with an estimated 20,000 people in the US living with hemophilia A. The condition is characterized by bleeding in joints, either spontaneously or in a provoked joint by trauma. Bleeding can occur in many different body areas as well (e.g., muscles, central nervous system). The bleeding manifestations can lead to substantial morbidity such as hemophilic arthropathy. Disease severity is usually defined by the plasma levels or activity of Factor VIII classified as follows: severe (levels < 1% of normal), moderate (levels 1% to 5% of normal), and mild (levels > 5% to < 40% of normal); phenotypic expression may vary. Approximately 50% of patients with hemophilia A are categorized as having severe disease.

Guidelines

Guidelines have not addressed Altuviiio. Guidelines for hemophilia from the National Hemophilia Foundation (March 2023)⁶ and the World Federation of Hemophilia (2020)⁷ recognize the important role of Factor VIII products and Hemlibra® (emicizumab-kxwh subcutaneous injection) in the management of hemophilia A in patients. The National Bleeding Disorders Foundation recognize Altuviiio as a product with a prolonged half-life.

Dosing Considerations

Dosing of clotting factor concentrates is highly individualized. The National Hemophilia Foundation's Medical and Scientific Advisory Council (MASAC) provides recommendations regarding doses of clotting factor concentrate in the home (2016).⁸ The number of required doses varies greatly and is dependent on the severity of the disorder and the prescribed regimen. Per MASAC guidance, patients on prophylaxis should also have a minimum of one major dose and two minor doses on hand for breakthrough bleeding in addition to the prophylactic doses used monthly. The guidance also notes that an adequate supply of clotting factor concentrate is needed to accommodate weekends and holidays. Therefore, maximum doses in this policy allow for prophylactic dosing plus three days of acute bleeding or perioperative management per 28 days. Doses exceeding this quantity will be reviewed on a case-by-case basis by a clinician.

Medical Necessity Criteria

Altuviiio is considered medically necessary when the following is met:

FDA-Approved Indication

1. Hemophilia A. Approve for 1 year if the patient meets ONE of the following (A or B):

A) Initial Therapy. Approve if the patient meets ALL of the following (i, ii, and iii):

- i.** Altuviiio is being used in at least ONE of the following scenarios (a, b, or c).
 - a)** Routine prophylaxis; OR
 - b)** On-demand treatment and control of bleeding episodes; OR
 - c)** Perioperative management of bleeding; AND
- ii.** Patient meets ONE of the following (a or b):
 - a)** Patient meets BOTH of the following [(1) and (2)]:
 - (1)** Factor VIII inhibitor testing has been performed within the past 30 days;
AND
 - (2)** Patient does not have a positive test for Factor VIII inhibitors ≥ 1.0 Bethesda units/mL; OR
 - b)** Patient has not received Factor VIII therapy in the past; AND

- iii. Medication is prescribed by or in consultation with a hemophilia specialist; OR
- B) Patient is Currently Receiving Altuviiio or Has Received Altuviiio in the Past.** Approve if the patient meets the ALL of following (i, ii, and iii):
 - i. Altuviiio is being used in at least ONE of the following scenarios (a, b, or c):
 - a) Routine prophylaxis; OR
 - b) On-demand treatment and control of bleeding episodes; OR
 - c) Perioperative management of bleeding; AND
 - ii. Patient meets ONE of the following (a or b):
 - a) Patient meets BOTH of the following (1 and 2):
 - (1) Factor VIII inhibitor testing has been performed within the past 365 days; AND
 - (2) Patient does not have a positive test for Factor VIII inhibitors ≥ 0.6 Bethesda units/mL; OR
 - b) According to the prescriber, patient does not have clinical manifestations suggesting the presence of Factor VIII inhibitors; AND
- Note: Inhibitors may be present if bleeding is not well controlled, there is decreased responsiveness to Factor VIII therapy, and/or if expected Factor VIII activity plasma levels are not achieved.
- iii. Medication is prescribed by or in consultation with a hemophilia specialist.

Dosing. Approve the following dosing regimens (A, B, and/or C):

- A) Routine prophylaxis:** approve up to 50 IU per kg intravenously no more frequently than once weekly; AND/OR
- B) On demand treatment and control of bleeding episodes:** approve up to 50 IU per kg intravenously with additional doses once every 2 to 3 days for up to 10 days per episode; AND/OR
- C) Perioperative management of bleeding:** approve up to 50 IU per kg intravenously and provide for additional doses once every 2 to 3 days for up to 10 days per procedure.

Altuviiio for any other use is considered not medically necessary. Criteria will be updated as new published data are available.

Coding Information

Note: 1) This list of codes may not be all-inclusive.

2) Deleted codes and codes which are not effective at the time the service is rendered may not be eligible for reimbursement.

Considered Medically Necessary when criteria in the applicable policy statements listed above are met:

HCPCS Codes	Description
J7214	Injection, Factor VIII/von Willebrand factor complex, recombinant (Altuviiio), per Factor VIII IU

References

1. Altuviiiio™ intravenous infusion [prescribing information]. Waltham, MA: Bioverativ/Sanofi; September 2024.
2. Mancuso ME, Mahlangu JN, Pipe SW. The changing treatment landscape in haemophilia: from standard half-life clotting factor concentrates to gene editing. *Lancet*. 2021; 397:630-640.
3. Croteau SE. Hemophilia A/B. *Hematol Oncol Clin North Am*. 2022;36(4):797-812.
4. Franchini M, Mannucci PM. The more recent history of hemophilia treatment. *Semin Thromb Hemost*. 2022;48(8):904-910.
5. Peyvandi F, Garagiola I, Young G. The past and future of haemophilia: diagnosis, treatments, and its complications. *Lancet*. 2016;388(10040):187-197.
6. National Bleeding Disorders Foundation. MASAC (Medical and Scientific Advisory Council) recommendations concerning products licensed for the treatment of hemophilia and selected disorders of the coagulation system (endorsed October 2, 2025). MASAC Document #290. Available at: <https://www.hemophilia.org/sites/default/files/document/files/MASAC-Products-Licensed.pdf>. Accessed on February 17, 2025.
7. National Hemophilia Foundation. MASAC (Medical and Scientific Advisory Council) recommendations regarding doses of clotting factor concentrate in the home (Revised June 7, 2016). MASAC Document #242. Adopted on September 3, 2020. Available at: <https://www.hemophilia.org/sites/default/files/document/files/242.pdf>. Accessed on February 19, 2025.

Revision Details

Type of Revision	Summary of Changes	Date
Annual Revision	Updated the scenarios for Altuviiiio use. Added Factor VIII testing requirements. Updated the specialist prescribing requirement. Added criteria for a patient currently receiving Altuviiiio or has received Altuviiiio in the past.	08/15/2024
Selected Revision	In Hemophilia A, for Initial therapy, the threshold for a positive inhibitor test was changed to ≥ 1.0 Bethesda units/mL; previously, it was ≥ 0.6 Bethesda units/mL. It was added that a patient who has not received Factor VIII therapy in the past is not required to meet the inhibitor testing requirements. For a Patient Currently Receiving Altuviiiio or has received Altuviiiio in the past, the Factor VIII inhibitor testing timeframe was changed to within the past 365 days; previously, the timeframe was within the last 30 days. The wording "prescribing physician" was replaced with "prescriber."	01/15/2025
Selected Revision	Updated CPT Coding: Removed: J7205 Added: J7214	01/15/2025
Annual Revision	No criteria changes.	5/15/2025

The policy effective date is in force until updated or retired.

"Cigna Companies" refers to operating subsidiaries of The Cigna Group. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life Insurance Company, Connecticut General Life Insurance Company, Evernorth Behavioral Health, Inc., Cigna Health Management, Inc., and HMO or service company subsidiaries of The Cigna Group. © 2025 The Cigna Group.